


**IN THE SPECIFICATION:**

Please substitute page 4, line 19 through page 6, line 3, with the following:

The at least one silicone copolymer results from the addition reaction, in the presence of a catalyst, of at least:

- (a) one polysiloxane of formula (II):



in which:

- $R_4$ , which may be identical or different, are independently chosen from groups that can react by chain addition reaction such as, for example, a hydrogen atom or aliphatic groups comprising an ethylenic unsaturation, such as vinyl, allyl and hexenyl groups;
- $R_5$  in formula (II), which may be identical or different, are independently chosen from hydroxyl, alkyl, alkenyl, cycloalkyl, aryl, and alkylaryl groups, and can optionally further comprise at least one functional group, such as at least one functional group chosen from ethers, amines, carboxyls, hydroxyls, thiols, esters, sulfonates and sulfates; wherein:
  - the alkyl groups comprise, for example, 1 to 20 carbon atoms; the alkenyl groups comprise, for example, from 2 to 10 carbon atoms; the cycloalkyl groups comprise, for example, 5 or 6 carbon atoms; the aryl

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
[www.finnegan.com](http://www.finnegan.com)

groups comprise, for example, phenyl groups; and the alkylaryl groups comprise, for example, from 7 to 20 carbon atoms;

- In one embodiment, R<sub>2</sub> is chosen from methyl;
- n is an integer wherein the polysiloxane of formula (II) has a kinematic viscosity ranging from 1 to 1 x 10<sup>6</sup> mm<sup>2</sup>/s, for example, n may range from 5 to 5000; and
- (b) at least one silicone compound comprising at least one and not more than two groups capable of reacting with the groups R<sub>4</sub> of the at least one polysiloxane of formula (II), wherein:
  - at least one of the at least one polysiloxane of formula (II) and the at least one silicone compound (b) comprises an aliphatic group, such as a C<sub>2</sub>-C<sub>6</sub> aliphatic group, comprising an ethylenic unsaturation.

The at least one silicone compound (b) can be another polysiloxane of formula (II) in which at least one and not more than two groups R<sub>4</sub> of the at least one silicone compound (b) can react with the groups R<sub>4</sub> of the at least one polysiloxane of formula (II).

**IN THE CLAIMS:**

Please cancel claim 93 without prejudice or disclaimer and replace claims 1, 72, 77, 81, 82, 84, and 85 with amended claims 1, 72, 77, 81, 82, 84, and 85, as follows:

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP  
  
1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

1. (Amended) A cosmetic composition comprising:  
(1) at least one conditioner chosen from:  
- synthetic oils;